



Physics 101 Lab Manual

Dr. W.A. Atkinson
Southern Illinois University, Carbondale
Physics Department
Updated by Dr. Foudil Latioui

January 23, 2012

Contents

| | |
|--|------------|
| How to Write Labs | vii |
| -1.1 What You Need to Bring to the Lab | vii |
| -1.2 The Pre-Lab | vii |
| -1.3 Structure of the Labs | vii |
| -1.4 How to Draw Graphs | viii |
| 0 Pre-Lab Questions | 1 |
| 0.1 Pre-Lab 1: Observations of Venus and Mars | 2 |
| 0.2 Pre-Lab 2: Diffusion | 4 |
| 0.3 Pre-Lab 3: Observations of Motion | 7 |
| 0.4 Pre-Lab 4: Conservation of Mechanical Energy | 11 |
| 0.5 Pre-Lab 5: Electricity | 14 |
| 0.6 Pre-Lab 7: Atomic Spectra | 16 |
| 0.7 Pre-Lab 8: Radioactivity | 18 |
| I Planets and Atoms | 19 |
| 1 Lab 1: Observations of Venus and Mars | 21 |
| 1.1 Objectives | 21 |
| 1.2 The Earth's Orbit | 21 |
| 1.3 Phases of the Moon | 22 |
| 1.4 The Phases and Orbital Motion of Venus | 23 |
| 1.5 Mars | 25 |
| 1.6 Summary & Conclusions | 25 |
| 2 Lab 2: Diffusion | 31 |
| 2.1 Objectives | 31 |
| 2.2 Diffusion of Ink in Water | 31 |
| 2.3 Microscopic Model | 33 |
| 2.4 Summary & Conclusions | 35 |

| | | |
|------------|---|-----------|
| II | Newton's Universe | 41 |
| 3 | Lab 3: Observations of Motion | 43 |
| 3.1 | Objectives | 43 |
| 3.2 | Motion on an Inclined Air Track | 43 |
| 3.3 | The Concept of Acceleration | 46 |
| 3.4 | Summary & Conclusions | 47 |
| 4 | Lab 4: Conservation of Mechanical Energy | 51 |
| 4.1 | Objectives | 51 |
| 4.2 | Definitions | 51 |
| 4.3 | Theory | 51 |
| 4.4 | General Procedure | 52 |
| 4.5 | Detailed Procedure | 53 |
| 4.6 | Analysis | 54 |
| 4.7 | Graphical analysis | 56 |
| 4.8 | Summary & Conclusion | 57 |
| III | The World of Electricity | 61 |
| 5 | Lab 5: Electricity | 63 |
| 5.1 | Objectives | 63 |
| 5.2 | Electrostatics | 63 |
| 5.3 | The Concept of "Ground" | 67 |
| 5.4 | Conductors and Insulators: | 68 |
| 5.5 | Simple Circuits: | 69 |
| 5.6 | Summary & Conclusions | 71 |
| IV | Within the Atom | 73 |
| 6 | Lab 6: Atomic Spectra | 75 |
| 6.1 | Objectives | 75 |
| 6.2 | The spectroscope and the visible spectrum | 75 |
| 6.3 | Spectra of Gases | 78 |
| 6.4 | Measuring Spectra | 79 |
| 6.4.1 | Calibrating the Spectroscope | 79 |
| 6.4.2 | Measuring Unknown Gases | 79 |
| 6.5 | Bohr's atomic model and Plack's constant | 80 |
| 6.6 | Summary & Conclusions | 81 |

| | |
|---|-----------|
| 7 Lab 7: Radioactivity | 83 |
| 7.1 Objective | 83 |
| 7.2 Radiation | 83 |
| 7.2.1 Kinds of radiation | 83 |
| 7.3 The Geiger Counter | 84 |
| 7.4 The Cloud Chamber | 87 |
| 7.4.1 Operating the cloud chamber | 87 |
| 7.4.2 Optimizing the operating conditions | 88 |
| 7.5 Summary & Conclusions | 89 |

How to Write Labs

-1.1 What You Need to Bring to the Lab

You will need to bring the following to the labs:

- Pencils and an eraser,
- A 30 cm (12 inch) ruler,
- A calculator,
- Loose-leaf paper on which to write the lab.

-1.2 The Pre-Lab

Each lab has a set of pre-laboratory exercises which must be completed *before* you arrive at the lab. Pre-lab questions will be graded along with the labs.

-1.3 Structure of the Labs

The first page of your lab should include

- the title
- your name and student number
- your lab partner's name

During the labs, you will be required to answer questions, draw sketches, make graphs and tables of data and write a short summary.

Questions are to be answered on loose-leaf paper, unless stated otherwise. They must be written in proper english. Please avoid point-form answers.

Some labs require simple **calculations** to be done. Remember to show how the calculation is done. If a calculation is repeated many times (for example, when filling out a table), the details only need to be shown once.




Figure 1: Sample graph showing how to draw a smooth curve.

For each lab, a short **summary** must be written. The summary should describe (a) the main purpose of the lab and (b) your most important findings. The purpose of the summary is to help you see the “big picture”, so you shouldn’t simply repeat all of your results. When you write the summary, make sure you look at the objectives for the lab, and say something about each of them. The summary should answer two questions: “What physical principle or phenomenon did we study?” and “What did we actually learn about it?” The summary must be written in proper english.

-1.4 How to Draw Graphs

A sample graph is shown in Fig. 3. When you draw a graph, pay attention to the following rules:

- All graphs need to have a *title* and a *number*. For example “Graph 1: Radiation penetration in lead”.

- When drawing a graph of “A” versus “B”, put “A” on the vertical axis and “B” on the horizontal. This is so that your graph shows how a measurement (on the vertical axis) changes at intervals of the horizontal axis (which are often intervals of time).
- Graphs should be drawn in pencil.
- Graphs must be drawn on graph paper.
- Make the graphs big. Use as much of the page as possible *BUT* keep it simple. The entire page could have been used in Fig. 3 if each square were 0.77 m instead of 1 m, but this just makes graphing complicated, and doesn’t add anything to the final product.
- Make sure you label the axes of your graph. If the x-axis indicates time measured in seconds, then it should read “time (s)”.
- Make sure you draw the scale on the graph.

When you are asked to draw a **smooth curve** through your data, the curve generally will not pass through all the data points. This is justified because there is always a small amount of uncertainty (known as **error**) in making a measurement.

If the graph is a straight line, then often we want to find the **slope** of the graph. The slope of a straight line is found by drawing a triangle like the one shown in Fig. 2, and determining the **rise** (the length of the triangle in the y -direction) and the **run** (the length of the triangle in the x -direction). The slope is then given by

$$\text{slope} = \frac{\text{rise}}{\text{run}}$$

Notice that if the curve goes down, then the rise is negative and the slope is negative. Notice also that the **units** of the slope are given by the units of the y -axis over the units of the x -axis. If the y -axis is cm, and the x -axis is seconds, then the slope of a line on the graph is in cm/s. Finally, remember to **show your calculations** on the graph, as done in Fig. 2.




Figure 2: Sample graph showing how to calculate the slope.




Figure 3: Sample graph for making plots.

Chapter 0

Pre-Lab Questions

0.1 Pre-Lab 1: Observations of Venus and Mars

Name _____ Section _____ Date _____

You may answer these questions directly on the figures.

This section looks at the simplest orbital system: the earth and the sun. The aim of this section is to give you some practice connecting what we see on earth to the earth's orbital motion. Figure 1 shows two views of the earth. Figure 1 (a) shows a "side-view", with North pointing upwards. Figure 1 (b) shows a "top-view", with North pointing out of the page towards you.

Question #:1 In Fig 1 (b) there are three arrows shown at point A on the earth's equator. The arrows are labelled 1, 2, and 3. For a person standing at point A, which arrow points East, which points West and which points straight up into space?

- Figure 2 is a schematic figure (not to scale) of the earth in orbit about the sun. As in Fig. 1 (b), the view is from the "top", so that North points directly out of the page towards you. The earth is shown at a particular instant in time.

Question #:2 The points A and B indicate two different points on the earth's surface. At each point, draw arrows indicating West and East (a total of four arrows), assuming that North points out of the page towards you. Next, using the fact that the sun rises in the East, draw the direction of the earth's daily rotation.

- The time at a particular point on the earth's surface is determined by it's location relative to the sun. If the sun is directly overhead, then it is noon. On the opposite side of the earth, where "up" points directly away from the sun, it is midnight. At points half-way in between midnight and noon, it is either 6 pm or 6 am.

Question #:3 Estimate the time to the nearest hour at points A and B.




Figure 1: Two perspectives of the earth. In (a), the four compass directions are also shown at a particular point on the earth's surface in the Northern hemisphere. In (b), the North pole points directly out of the page towards you [ie. you are looking down from the "top" of (a)].




Figure 2: The earth in orbit around the sun.

0.2 Pre-Lab 2: Diffusion

Name _____ Section _____ Date _____

Reading: Hobson Ch.2, section 2.1 & section 2.2
Try to understand the comment of fig.2.1 of the textbook.
Review section 2.3 of the lab on diffusion

Question #:1 Define diffusion. *Hint:* Look it up in Wikipedia, the free online encyclopedia. use key words such as *molecular diffusion*, *Brownian motion*

- Figure 3 shows a cartoon of a random walker. Initially, the walker starts at 0. A coin is tossed and the walker moves one position to the right if the coin lands “heads” and one position to the left if the coin lands “tails”. We represent the position of the walker by a number r .
- Perform your own random walker experiment. Start at $r = 0$ and toss a coin 20 times. For each “tails” subtract 1 from r , and for each “heads” add 1 to r . Record this in a table. Note that r can be positive or negative.
- Read the section “How to draw graphs” in the introduction to the lab manual. Draw a graph of r vs. N (the number of coin tosses), following the guidelines set out in the introduction.




Figure 3: The random walker. If the coin-toss is “heads” the walker moves one step to the right; if the coin-toss is “tails” the walker moves one step to the left. In this figure, the walker is moving from $r = 0$ to $r = 1$.




Figure 4: Average distance travelled for many random walkers. This data is generated by having a computer toss a coin 100 times for each random walker, and then averaging the results.

Question #2 Describe, in words, the relationship between r and N for your random-walker experiment. In other words, is there a pattern in the way r changes when you increase N ? Is there anything predictable about the motion of your random walker?

- Figure 4 shows the average of $|r|$ for 10 random walkers and for 1000 random walkers. The symbol $|r|$ refers to the *absolute value* of r , which means dropping the minus sign (if there is one) in front of the number. Thus $|-5| = 5$, $|-3.7| = 3.7$, and $|10| = 10$. The figure shows the average distance of 1000 random walkers from their starting place after 1, 2, etc. coin tosses. This data was generated by a computer which tossed a coin 100 times for each random walker. The data was then averaged over all the random walkers. The data shows that, for 1000 walkers, the random walkers have moved an average of 1.87 steps away from their origin after 5 coin tosses. This means that some walkers have moved 1 step away, some have come back where they started from, and a few have even moved a full 5 steps away.

Question #3 Compare the average motion of the random walkers with your coin tossing experiment. What do you notice about the *average* motion of the random walkers as the number of walkers is increased? Note that each *individual* random walker still moves completely randomly!

Question #4 From your graph, can you estimate what $|r|$ will be for your single random walker after 30 coin tosses? How much confidence do you have in your estimate?

Question #5 From Fig. 4 estimate the average value of $|r|$ for 1000 random-walkers after 150 coin tosses. How much confidence do you have in your estimate?

Question #6 Which has more predictability, the motion of an *individual* random walker or the *collective* (average) motion of a large number of random walkers?

- A theory or an equation has predictive power if it can predict the outcome of an experiment which has not yet been performed.

0.3 Pre-Lab 3: Observations of Motion

Name _____ Section _____ Date _____

Reading: Hobson Ch.3: section 3.4 & section 3.5

Galileo found that gravity causes constant acceleration along an incline. Assuming an object starts from rest:

- The relationship between distance (s), acceleration (a), and time (t) is: $\Delta s = \frac{1}{2}at^2$, where Δ refers to the fact that the distance measurement measures the difference between its position at the beginning and end of the experiment.
- The relationship between speed, acceleration and time is: $v = at$
- Keep in mind that A^2 (A squared) is A multiplied by A, not A multiplied by 2.

Question #:1

- A cart on a linear track has a uniform acceleration of $0.172m/s^2$. What is the velocity of the cart 4.00 s after it is released from rest?
- How far does this cart travel during these 4 s?
- From the first equation, give the expression of the acceleration a

In table 1 is a data set of time and distance traveled.

Question #:2

- Complete table 1
- find the average acceleration

| | | | | | | | |
|---------------|------|------|------|------|------|------|------|
| $\Delta s(m)$ | 0 | 0.25 | 0.50 | 0.75 | 1.00 | 1.25 | 1.50 |
| $t(s)$ | 0.00 | 0.45 | 0.71 | 0.77 | 0.98 | 1.07 | 1.12 |
| t^2 | | | | | | | |
| a | | | | | | | |

Table 1: Average values of time are below the corresponding values of distance

Question #3 use the collected data in table 1 and:

- Plot distance versus time
- is this a uniform motion? Explain

Question #4 use the collected data in table 1 and:

- Plot $2\Delta s$ versus t^2
- Give an interpretation of the slope of this graph
- Extract the average acceleration from the graph. and compare with the previously calculated one.




Figure 5: Distance Versus Time.




Figure 6: Twice the distance versus Time squared

0.4 Pre-Lab 4: Conservation of Mechanical Energy

Name _____ Section _____ Date _____

Reading: Hobson Ch.6: section 6.5 & section 6.6

- Potential energy is given by $PE = mgh$
- Kinetic energy is given by $KE = \frac{mv^2}{2}$




Figure 7: Trade in joules between potential and kinetic energy

Consider the situation in figure 7, the total mechanical energy is 6 J. There is no loss or gain of mechanical energy, only a transformation from kinetic energy to potential energy (and vice versa). The ball, of mass $m = 1\text{kg}$ is released from rest. Answer the following questions:

Question #:1 At point A:

- What is the potential energy of the ball?
- What is the height of the ball?
- What is the velocity of the ball at that point?
- What is the kinetic energy?

Question #:2 At point B, if the potential energy is equal to the kinetic energy ($PE = KE = 3J$)

- What is the height H of the ball?
- What is the velocity of the ball at that height?

Question #:3 At point C, the potential energy is zero.

- What is the kinetic energy of the ball?
- What is the velocity of the ball at that point?
- How would you describe the velocity at this particular point? Explain

Question #:4 At point D, the ball is at the same height as in point A, it reaches a maximum height then stops before reversing direction.

- What is the potential energy?
- What is the kinetic energy of the ball?
- What happened to kinetic energy at that position?

Question #:5 While the pendulum is oscillating between positions, use the words **increase**, **decrease**, or **constant** to describe the trend of the different forms of energies. Complete table 2

| | Potential Energy | Kinetic Energy | Mechanical Energy |
|-------------|------------------|----------------|-------------------|
| From A to C | | | |
| From C to D | | | |

Table 2: Trend of the different forms of energy of a swinging pendulum

Question #:6

- Using figure 7, identify the corresponding positions A , B and C in figure 8
- Discuss the different energies, and speeds of the glider on those points based on your study of the pendulum.




Figure 8: Sliding glider down an incline

0.5 Pre-Lab 5: Electricity

Name _____ Section _____ Date _____

Reading: Hobson Ch.8-section 8.1 and 8.3

Use the principles given in the Objectives part of the lab (section 5.1) to answer the following questions:

Question #:1 Look at Fig. 9. Two charges (labelled A and B) are placed in some object (for example, a piece of glass or a piece of metal). Assume for the moment that both charges are electrons. If the material is a conductor, how will the electrostatic forces make them move? Where will they stop?

Question #:2 How will the charges move in the conductor if charge A is an electron and charge B is a positive ion? Where will they stop?




Figure 9: Two charges (A and B) are placed in an object. A third charge (C) is added later.

Question #:3 After the charges in the previous problem stop moving, a third charge (charge C) is put into the conductor. Do you expect it to feel a force? Give your reasoning.




Figure 10: Some extra negative charge is placed in a conductor. How does it move?

Question #4 How will the charges in Fig. 9 move due to the electrostatic forces if the material in which they are placed is an insulator?

Question #5 Figure 10 shows a conductor that has some extra electrons placed on it. Describe how the excess negative charge moves? How is it distributed when it stops moving?

0.6 Pre-Lab 7: Atomic Spectra

Name _____ Section _____ Date _____

Reading: Hobson Ch.9.3, 9.4, 9.6, 13.6 & 13.7

The following hypothesis is a *classical theory* which tries to explain why a light bulb emits light. It is classical because it ignores the wave-like properties of atoms.

Hypothesis 1 *Incandescent light comes from a tungsten filament encased in a glass bulb.*

- 1. Electric current passes through the tungsten filament and heats it up.*
- 2. Tungsten atoms vibrate randomly when heated. They move more when they are hotter.*

Question #:1 Use the fact that atoms are made of charged protons and electrons to explain why the random vibration produces light.

Question #:2 Consider an experiment where you produce waves on a string by moving one end up and down with your hand. What is the relationship between the frequency of your hand (ie. the number of times per second you move your hand up and down) and the frequency of the wave on the string?

Question #:3 What do you think is the relationship between the frequency of atomic vibration in the tungsten filament and the frequency of emitted light? Based on this hypothesis, what frequencies should be present in the light emitted by the filament?

Question #:4 If the atoms are hotter, then we can expect that they will move *faster* and *farther*. What do you think that this would do to the brightness of the light? What about the color of the light?

Question #:5 What is the energy transformation which occurs when light is produced? In other words, where does the energy for the light come from? Give answers in terms of both *macroscopic* and *microscopic* energies.

0.7 Pre-Lab 8: Radioactivity

Name _____ Section _____ Date _____

Reading: Lab Manual section 7.2, Hobson Ch. 14.1-14.3

Question #:1 What are the four fundamental forces?

Question #:2 Which fundamental force holds the nucleus together?

Question #:3 Which fundamental force is responsible for α -decay?

Question #:4 Which fundamental force is responsible for β -decay?

Question #:5 Write down the reaction equation for the β -decay of ^{90}Sr (Strontium-90).

Question #:6 Write down the reaction equation for the α -decay of ^{239}Pu (Plutonium-239).

Part I

Planets and Atoms

Chapter 1

Lab 1: Observations of Venus and Mars

Name _____ Section _____ Date _____

1.1 Objectives

1. To learn what information can be deduced about the planets based on changes in their appearance when they are observed over a period of several months.
2. To understand differences in the appearances of superior and inferior planets.

1.2 The Earth's Orbit

The following questions are based on Fig. 1.2. The questions may be answered directly on the figure.

Question #:1 If Fig. 1.2 shows the position of the earth on February 13, draw, as accurately as possible, the position of the earth on April 13, two months later. Recall that the earth orbits the sun in the same direction that it rotates on its axis. The angular distance travelled by the earth can be computed. Recall that the earth travels 360 degrees in one year, so that in 19 days it travels

$$\frac{19}{365} \times 360 = 18.7 \approx 19 \text{ degrees.}$$

From this we can see the earth travels approximately one degree per day. You may use this approximation throughout this lab.

Question #2 Draw the new position of point A on April 13, if it is the same time of day as shown for February 13.

Question #3 A star is directly overhead at midnight on February 13. Draw an arrow showing the direction to the star. At what time of day will the star be directly overhead on April 13? You can figure this out from the sketch you drew in Question 2. (see following note as a hint)

- Note that stars are very far away, so that the arrows pointing to the star on February 13 and April 13 will be nearly parallel. For the scale shown in Fig. 1.2 (representing the earth's orbit by a circle a few inches across), the nearest stars would be approximately 1 mile away.

1.3 Phases of the Moon

The “phase” of the moon refers to how it is illuminated by the sun. The “full moon” is the phase in which the entire side of the moon facing the earth is illuminated. A “crescent moon” has only a small sliver illuminated. In this section, you will examine the relationship between the phases of the moon, and the orbital motion of the moon.

Figure 1.3 shows the moon in orbit around the earth. The moon is shown at three different points in its 28-day orbit. The following questions may be answered on the figure.

Question #4 Assuming that North points out of the page, draw the direction of rotation of the earth, and the direction the moon travels along its orbit. The moon orbits in the same direction the earth spins.

Question #5 At each of the three locations shown, shade the night side of the moon with a pencil. Note that light comes as parallel rays from the sun, as shown in the figure.

Question #6 For each case, make a sketch of the moon showing how you would see it from Earth.

Question #:7 Indicate which of the three phases of the moon is closest to “full” and which is “crescent”.

1.4 The Phases and Orbital Motion of Venus

In this section, you will make the connection between a sequence of telescope observations of venus and venus’s orbital motion. Venus is an *inferior* planet, which means that the radius of its orbit is smaller than that of the earth. This section is a little more complicated than the previous sections because the motions of both venus *and* the earth need to be considered.

You will be given a data sheet showing a series of telescope-photos of venus taken over a 5 month period. Answer the following questions regarding the data.

Question #:8 Figure 1.1 shows two possible arrangements of the earth-sun-venus system. Which of these corresponds most closely to the January 19 observation? Give your reasons.

Question #:9 Why does venus change size in the photos?

Question #:10 Using the fact that North is up in the photos, how can you tell that the observation on Jan. 19 was made in the evening (and not just before dawn)?

Question #:11 Figure 1.4 shows the earth-sun-venus system with the earth’s position shown on the date of the first venus photo (Jan. 19/01). Use the data shown in the handout to find the location of venus on the observation dates listed below. This must be done in several steps.

- a.) First you need to determine where the earth is on the observation dates. Make a table like the one shown below




Figure 1.1: The earth-sun-venus system.

| Date | Days Elapsed since Jan. 19 |
|---------|-------------------------------|
| Jan. 19 | 0 |
| Feb. 20 | 32 |
| Mar. 21 | |
| Apr. 13 | |
| May 8 | |

The left-hand column shows the dates on which the observations were made. The right-hand column shows how much time has elapsed since the Jan. 19 observation. For example, Feb. 20 is

$$20 + (31 - 19) = 32 \text{ days}$$

later than Jan. 19. This is the number of days in February (20), added to the number of days left in January, which is the total number of days in January (31) minus the number which have already happened, thus $31-19$. Since we are looking for the days elapsed since January 19th, you also have to add the number of days already elapsed from the previous months until January 19th, in this case zero (though it won't be the next month). Recall that the earth travels one degree in one day. Complete the table for the remaining observation dates.

- b.) Draw, directly on fig. 1.4 and as accurately as possible, the positions of the earth on each of the observation dates.
- c.) Next, draw as accurately as possible the positions of venus on the observation dates. To do this, you will need to use information contained in the handouts; the telescope observations tell you how the size and illumination (by the sun) of venus changes. From this, you can determine the relative positions of the earth and venus.
- d.) Based on your results, estimate (very roughly) the length of the venusian year.

1.5 Mars

You have also been given a series of photos of mars taken from the Hubble Space Telescope over a period of nearly 1 year. Mars is a *superior* planet, which means that the radius of its orbit is larger than that of the earth.

Question #:12 Compare the martian and venusian photos. Why do we sometimes see a “crescent venus” (only a thin sliver is illuminated) but never a “crescent mars”? Note that the radius of mars’s orbit is *bigger* than the radius of earth’s orbit.

Question #:13 Why can you often see mars at midnight, but never venus?

1.6 Summary & Conclusions

Write a short summary of the lab which addresses the issues raised in the Objectives. For example, you might consider the following questions:

- How do the appearances of venus and mars change as they move?
- Why do they change appearance?
- In what ways are they similar? Different?
- What is the reason for their similarities/differences?

Question #:14 [Bonus Question] Are the data sheets (the photos of venus and mars) compatible or incompatible with a geocentric (earth-centered) model of the solar system? Is there any feature of the data which can be used to rule a geocentric model out?




Figure 1.2: The earth in orbit around the sun.





Figure 1.3: The moon in orbit around the earth.



Earth: 01/19/01

Figure 1.4: The earth-sun-venus system.




Figure 1.5: The earth-sun-mars system.

Chapter 2

Lab 2: Diffusion

Name _____ Section _____ Date _____

2.1 Objectives

This lab studies several important concepts:

1. The concept of predictive power.
2. The microscopic model of diffusion.
3. The difference between the microscopic and macroscopic motion of ink molecules in water.

2.2 Diffusion of Ink in Water

- Fill the bottom of the petrie dish with a thin layer of water. Read the instructions below thoroughly before beginning your experiment.
- Set the petrie dish on top of the ruler so that the centimeter scale of the ruler is clearly visible and passes through the center of the dish, as shown in Figure 2.1.
- Make a table like the one below:

| Time (s) | Position of left edge (cm) | Position of right edge (cm) | d (cm) |
|----------|----------------------------|-----------------------------|----------|
| | | | |




Figure 2.1: Experimental Setup for Diffusion Lab.

- In the first three columns you will record the position of the left and right edges of the ink drop at different times as it expands. List the times before you start based on the instructions below. In the fourth column you will record the diameter d of the ink-drop.
- Carefully place a *small* ink drop near the middle of the petrie dish, over the edge of the ruler (as shown in Figure 2.1) and start timing. You need to centre the ink drop on the edge of the ruler so that you measure its diameter.
 - If you don't get a nice compact ink drop, you should start again. It may take several tries.
 - This experiment works best if the water is at room temperature. If the water is too cold, you will get convection currents.
- Use the clock on the wall or your cellphone to time the experiment. Make measurements of the positions of the left and right edges of the ink-drop at 15 second intervals for the first two minutes. After two minutes, make observations every 30 seconds. Continue until at least 180 seconds (3 minutes) have passed.
- Make a graph of the ink-drop diameter d vs. the elapsed time. Instructions on how to draw graphs are given in the introduction. Draw a smooth curve through the data points.

Question #:1 Describe in words the relationship between d and t . Is it predictable?




Figure 2.2: The path of an ink molecule (the large gray molecule) through water (the small black molecule).

Question #2 The ink-spot expands more slowly at large times. Does this mean that the individual molecules are slowing down?

2.3 Microscopic Model

Consider a simple model for diffusion, illustrated in Fig. 2.2. This is the *microscopic diffusion model*.

1. The Petrie dish contains two types of molecules—ink molecules and water molecules. The molecules move in random directions.
2. Ink molecules move in a straight line until they bump into another molecule.
3. During a collision, a molecule is scattered in a random direction. (In fact, the direction *isn't* really random. If we knew the details of exactly how each collision occurred then the direction of the molecules after each collision could be predicted exactly.)

4. The molecule continues moving in a straight line until it bumps into another molecule.
5. Because of this random scattering the ink molecules move in a random walk. This means that the motion of an individual molecule is unpredictable.
6. Even though the motion of each individual molecule is random, the collective motion (ie. the average motion of many ink molecules) is predictable. This collective motion is called diffusion.
7. The *rate* at which molecules diffuse is measured by the “Diffusion constant” D . A large value of D means that molecules spread out quickly.
8. The diffusion constant D is determined by the kind of molecules diffusing, by the medium through which they diffuse, and by the temperature.

Question #:3 The speed of a molecule is determined by its temperature (hot molecules move faster). How will temperature affect the diffusion constant D ? Why?

Question #:4 Since molecules are farther apart in gases, they travel farther between collisions. Will D be larger or smaller in gases than in liquids? Why?

- If diffusing molecules can be thought of as random-walkers, then it should be possible to make a connection between the microscopic diffusion model and the coin tossing experiment.
- The following questions try to show that the coin-tossing experiment is a useful “toy-model” for diffusion. That means it is a simplified model which contains the basic ingredients to explain diffusion.

Question #:5 Assume that the coin-tossing experiment can explain diffusion. What element of the microscopic diffusion model does the random-walker itself represent?

Question #:6 What microscopic property of the ink molecules does the variable r from the coin tossing experiment represent?

Question #:7 What element of the microscopic diffusion model does the variable N from the coin tossing experiment represent?

Question #:8 What does each coin toss represent?

Question #:9 What *macroscopic* property of the ink spot does the *average* r represent?

2.4 Summary & Conclusions


What have we done here? This lab has many pieces to it so it can be a bit confusing. We studied a “random walker” model of diffusing molecules. In order to check whether this model works, we made a prediction that we could test. In this case, the prediction had to do with how the size of the ink spot changes with time.


Question #:10 Compare your graph of d vs. t with the data for the average $|r|$ vs. N shown in Fig. 4. Are these two graphs consistent with each other? In other words, does your graph of d vs. t support or contradict the random walker model of diffusion?


Question #:11 Does this experiment prove that molecules exist? Does it support the theory that molecules exist? Justify your answer.

- Now write a brief discussion of the lab based on the objectives. You can turn each objective into a question by starting it with “What is...”

Question #:12 Bonus Activity: If you perform the ink spot experiment with cold water (the colder the better), then after a minute or two of observing, you should see *convection cells* form. Describe what you see, especially as it is different from when you did the experiment with warmer water. Draw pictures if it will help.







Part II

Newton's Universe

Chapter 3

Lab 3: Observations of Motion

Name _____ Section _____ Date _____

3.1 Objectives

- To study the motion of a glider on incline
- To understand the concept of acceleration

3.2 Motion on an Inclined Air Track

In this section, you will study the motion of a body accelerating under gravity. These experiments are very similar to ones performed by Galileo in the 17th century. He studied the motion of rolling balls on an inclined plane, and deduced, among other things, that falling bodies accelerate uniformly.

This lab examines acceleration more closely. You will look at the motion of a glider on an inclined air track. Your TA will demonstrate the use of the air track and photogate with the accessory photogate.

- First, level the air track. You can do this by adjusting the threaded legs on the track. When the air is on, the glider should remain stationary when you release it.
- Now, elevate the air track by turning the adjustable legs through 5 complete turns. (The end marked “140 cm” should be higher than the end marked “0 cm”).
- Make a table like the one below




Figure 3.1: Schematic diagram of the air track.

**Table 1: Motion on an air track
elevated by 5 turns**

| Position s_1 (cm) | Position s_2 (cm) | Distance Δs (cm) | Time t_1 (s) | Time t_2 (s) | Time t_3 (s) | average time t (s) |
|------------------------|------------------------|-----------------------------|-------------------|-------------------|-------------------|-------------------------|
| | | | | | | |
| 10 | 20 | | | | | |
| 10 | 30 | | | | | |
| 10 | 40 | | | | | |
| 10 | 60 | | | | | |
| 10 | 80 | | | | | |
| 10 | 130 | | | | | |

Here s_1 is the initial position of the glider *measured at the leading edge* of the glider, and t_1 , t_2 and t_3 are the times required for the glider to travel to the end of the inclined plane on three different trials.

- Place leading edge of the the flag on top of the glider just before the 10 cm mark (at s_1) on the inclined plane, along with the accessory photogate. Make sure the switch on the photogate is set to pulse mode and the other switch is set to the on position. Release the glider just before the position that it will trip the photogate, without giving it any sort of a push forward or backward. The photogate should time how long it takes to travel to the other photogate, positioned at s_2 on the air track. Record your answer in the table in the column labelled t_1 . Repeat the process twice more and record the answers under columns t_2 and t_3 . Record the average of t_1 , t_2 , and t_3 on the chart.
- Calculate $\Delta s = s_1 - s_2$, which is the total distance travelled by the glider. Record the value in the column marked 'Distance Δs '

Question #:1 List at least 2 important sources of error. In other words, give two reasons why t_1 , t_2 and t_3 differ from each other.

- Using the graph paper provided, carefully make a graph of Δs versus t , using the average t . Draw a *smooth* curve through the data points (remember, you don't need to hit each one, just give a good idea of the trend with your line). When you draw the curve, remember that when $\Delta s = 0$ cm, $t = 0$ s. A neat graph will make some of the following questions easier.

Question #:2 Based on your graph, does the glider speed up, slow down, or travel at a constant speed as it moves along the track? Hint: Estimate how far the glider moves in the first second, and in the last second.

Question #:3 What is the *average* speed of the glider during the last run (starting at $s_1 = 130$ cm)? Don't forget to show your calculation. Recall that average speed is equal to the distance divided by the total time.

- You have seen in class that the *instantaneous* speed can be found from the slope of the Δs versus t graph. Pick three (approximately) evenly-spaced values of t on your graph which cover the whole range of times measured in the experiment (that is, pick points at the beginning, middle and end).
- At those three points from the graph, draw a *tangent line* to the smooth curve you drew through your data. A tangent line is a line which grazes against a curved line, touching it but coming out on the same side of the curve on both directions before any intersection.
- Find the slope of the tangent line at each selected time. Instructions for finding slope are found in the Introduction to the lab manual. You will use points on the tangent line. The points you use to find the slope which are not on your smooth curve will not be used after this. This slope has units of cm/s and is the speed of the glider at that instant in time.

- Record your results for each tangent line in the table below:

Table 2: Speed of the glider down the incline

| Time t in seconds | Distance Δs in cm | Speed v in cm/s |
|---------------------|---------------------------|---------------------|
| | | |

Question #:4 Compare the instantaneous speeds you calculated with the average speed of the glider you calculated earlier. (for $s_1 = 130cm$)

3.3 The Concept of Acceleration

- We can define *acceleration* to mean that the speed of the glider increases as it moves down the track. The speed of the glider changes with respect to position and time. If you think about it for a minute, you will find that this is ambiguous. Galileo had to struggle with the idea of acceleration.

Definition: The average acceleration over some time interval $t_f - t_i$ is given by

$$a = \frac{v_f - v_i}{t_f - t_i}$$

where $v_f - v_i$ is the change in speed over the interval. Here, t_i and t_f are the beginning and ending points of the time interval, and v_f and v_i are the speed of the glider at t_f and t_i .

- Using the data you recorded in Table 2, make a graph showing the speed of the air glider versus the total time t the glider has moved.
- Draw a smooth curve through your data.

Question #:5 Describe the relationship between the speed of the glider and the time it travels. Is there a predictable pattern? Can you predict the speed of the glider after 30 s?

- From the definition, we can define an *instantaneous* acceleration by making the distance interval $t_f - t_i$ very small. This is equivalent to finding the *tangent* to the curve.
- Plot the three points from table 2 in a graph of speed vs time. Does the graph look linear (close to a straight line)? Find the slope of a line between the first and last points.

Question #:6 What are the units of acceleration (based on the definition earlier)?

Question #:7 Does the acceleration change as the glider moves down the air track? Explain.

Question #:8 From a series of careful experiments, Galileo determined that the acceleration of a body down an inclined plane is *uniform* (constant) provided he used Definition 2 for acceleration. Do your results support this? Do you find any significant deviations from Galileo's theory? ("Significant" means you can't simply blame them on human error in the timing.)

Question #:9 If you change the elevation (ie. the steepness) of the air track, how do you think that this will change the relationship between v and t . Will the shapes of the curves change? If so, how?

3.4 Summary & Conclusions

Summarize your main findings in this lab. What did you set out to investigate and what did you find? Remember to address the objectives for the lab.




Figure 3.2: Graph of distance traveled versus the average time: Δs Versus t_{av}




Figure 3.3: Speed of the glider Versus the time it travels down the incline. v Versus t

Chapter 4

Lab4:Conservation of Mechanical Energy

Name _____ Section _____ Date _____

4.1 Objectives

In this laboratory experiment, we are going to illustrate the conservation of mechanical energy:

- Quantitatively: through measurement and data analysis
- Qualitatively: through graphical analysis

4.2 Definitions

- Gravitational Potential Energy is an energy by virtue of **position**: $PE = mgh$
- Kinetic Energy is an energy of **motion**: $KE = \frac{mv^2}{2}$
- Mechanical Energy: $ME = KE + PE \Rightarrow ME = \frac{mv^2}{2} + mgh$

4.3 Theory

The Total Mechanical Energy of an object remains constant in the absence of nonconservative forces like friction. The conservation of energy is a fundamental tenet of science and has extremely broad applications in all technical fields. The two types of energy that will be under consideration in this laboratory experiment are the gravitational potential energy PE and the kinetic energy KE. The object in this case will be a glider on an air track system. There are two forces at work in this experiment - gravity and friction.

The frictional forces have been minimized by the use of the air track system and will therefore be neglected in our analysis. Gravity exerts a force on the glider and will contribute a gravitational potential energy (PE) to the Total Mechanical Energy. The normal force of the track pushing back on the glider is perpendicular to the direction of the glider motion and so will produce no work and will not affect ME.

Therefore, in this experiment the Total Mechanical Energy consists only of the Kinetic Energy (KE) and the Gravitational Potential Energy (PE). Since there are no non-conservative forces acting on the glider its Total Mechanical Energy is conserved.

4.4 General Procedure

- As the glider accelerates down the sloped air track you will make measurements of the glider's PE and KE for six different locations along the air-track.
- These six positions are located at 20 cm, 40 cm, 60 cm, 80 cm, 100cm and 120cm marks along the length of the track.
- The glider's PE can be calculated if we know its vertical height from the reference level of zero gravitational potential energy. We will choose the zero reference level to be the height of the sixth (bottom) y_6 which corresponds to 120cm-mark on the air-track.
- We will first measure the heights of each of the measurement positions relative to the surface of the table and record these in the first column of your Data Table 4.1.
- Then subtract off the height that you measured for the sixth (bottom) position y_6 from all six of the measured height values.
- Record these results in the second column of your Data Table 4.2.
- The velocity needed for calculating the glider's KE will be measured at five locations along the air track using the photogate. The TA will demonstrate how to use the photogate.
- The velocity of the glider, v , at each measurement location will be determined by the photogate which will time the passage of a narrow flag (a piece of paper or notecard carefully folded multiple times), mounted in the front of the glider as it passes through the photogate.
- Since the glider is at rest at the first location y_1 , the KE at that position is zero.
- After you place the photogate at one of the measurement locations and let the glider accelerate down the air track the photogate will provide the timing information that together with the flag length will enable you to calculate the glider's average velocity at the measurement location:

$$v = \frac{L}{\Delta t}$$
- After each velocity is calculated, record it in your Data Table 4.1.
- Once all the velocity-positions data have been collected, you will calculate the values of the three quantities, PE , KE , and $ME = KE + PE$ and record them in your Data table 4.2.




Figure 4.1: glider-air track system: heights y_i & positions along the incline

- Then you will plot each of these three variables versus position of the glider along the air track.
- These plots will allow you to test the validity of the claim that Mechanical Energy, as applied to our simple glider and air-track system, is conserved.

4.5 Detailed Procedure

- For better results, all of your observations should be recorded to 3 significant figures. You should carry 3 significant figures (three digits) in all of your calculations as well.
- Measure the mass of your glider using the available scale and record here $M = \text{----- } g, \text{----- } kg$
- Measure the length of the flag of the glider using the provided Caliper and record the length here $L = \text{---- } mm, \text{----- } m$.
- (Optional) Place a jack or a block of wood under the single leg of the air track. Set up the air-track with an inclination roughly similar to that shown in figure 4.1
- The initial vertical height measurements are denoted by the variable y_i . The measurements should be made using the calipers from the bottom of the air track to the table top at positions located at 20cm, 40cm, 60cm, 80cm, 100cm, and 120cm along the air track.
- The launching point should be the same (position 20 cm on the airtrack) for all trials.
- We will designate the height y_6 as the zero reference level of the gravitational PE .
- The difference in heights relative to this zero level will be designated by the variable h defined as $h_i = y_i - y_6$.

- Measure y_1 through y_6 , as shown in figure 4.1, to three significant figures. Calculate the h_i values and record them in your Data Tables 4.1 and 4.2
- Place the photogate at the first data collection point, $s = 40\text{cm}$. Switch the photogate to **GATE** mode. Make sure to **RESET** gate before each time you take data.
- The photogate should be rotated slightly about its mounting axis so that its body is perpendicular to the plane of the air track. The flag should not strike anything as the glider travels down the air track. The flag should be flush with the forward edge of the cart. After taking data at this point this alignment procedure will be repeated at each of the other positions
- Turn on the air supply to the air track. The glider will float on a cushion of air.
- Let the glider accelerate from its rest position at position 20cm mark.
- Record the velocity measured, when the glider passes through the photogate, in the Data Table 4.1. Repeat this velocity measurement two more times for a total of three velocity measurements at each photogate position.
- Repeat these velocity measurements with the photogate positioned at the remaining locations, and record the values in the corresponding Data Tables 4.1.
- Calculate the average velocity for each measurement location and record the results in both tables.

Complete table 4.1 remember that $h_i = y_i - y_6$.

Complete table 4.2. Assume $g = 9.8\text{m/s}^2$

4.6 Analysis

After completely filling in the two data tables, you are ready to answer questions and make your graphs.

Question #:1

- What is the gravitational potential energy of the glider at the very top (y_1 position)? Show your calculation in the space below

- Is it maximum or minimum?
- How is it compare to the total mechanical energy?

Question #:2

- At the height y_3 , what fraction of the total mechanical energy is gravitational potential energy?
Show your work.

- Could you have qualitatively predicted such an outcome? Explain.

- What will happen to the potential energy as the glider slides downward?

Question #:3

- What is the kinetic energy of the glider at the very bottom (y_6 position)?
Show your calculation in the space below

- Is it maximum or minimum?
- How is it compare to the total mechanical energy?

- what is the potential energy at this location?

- What happened to the potential energy?

Question #:4

- In this lab we neglected friction. If friction was present what would it do to the values of the velocities?

- How would this affect the kinetic energy values?

- What would this do to the value of the Total Mechanical Energy (ME)?

- Make a general statement about the effect of friction on a given system. (will it add or remove energy)

4.7 Graphical analysis

In this section, we will discuss the conservation of mechanical energy graphically. The vertical scale will be used to represent energy in joules and the horizontal scale will be used to represent the positions in centimeters. Use the data collected in table 4.2. Plot everything on the same graph.

1. Plot of potential energy versus position (PE Vs s):
 - Draw a smooth curve or line through the data
 - Describe the trend of the curve (increasing, decreasing or constant)

2. Plot of kinetic energy Versus position (KE Vs s):
 - Draw a smooth curve or line through the data.

- Describe the trend of the curve (increasing, decreasing or constant)
3. Plot of mechanical energy versus position(ME Vs s):
- Draw a smooth curve or a line through the data.
 - Describe the trend of the curve (increasing, decreasing or constant)
 - How can you tell from the graph that $PE + KE = Constant$

4.8 Summary & Conclusion

Question #:5

- Within reasonable limits of this experiment, were you able to test the validity of the following claim: *"in the absence of non-conservative forces(friction), the mechanical energy for the glider on the air track is conserved"*. Explain.
-
- "Energy is generally conserved. Mechanical energy is conserved when frictional forces are minimized." True or False?




Figure 4.2: Kinetic, Potential & Mechanical Energy Versus Position

| Position: s_i | Height: h_i | Repeated measurement of time: t_i | average velocities v_i |
|-----------------|---------------|---|--|
| 20 cm | $h_1 =$ | $t_1 =$ $t_2 =$ $t_3 =$ $t_{av} = \frac{t_1 + t_2 + t_3}{3} =$ | $v_{av1} = \frac{\text{flag length}}{t_{av}}$ $v_{av1} =$ |
| 40 cm | $h_2 =$ | $t_1 =$ $t_2 =$ $t_3 =$ $t_{av} = \frac{t_1 + t_2 + t_3}{3} =$ | $v_{av2} =$ |
| 60 cm | $h_3 =$ | $t_1 =$ $t_2 =$ $t_3 =$ $t_{av} = \frac{t_1 + t_2 + t_3}{3} =$ | $v_{av3} =$ |
| 80 cm | $h_4 =$ | $t_1 =$ $t_2 =$ $t_3 =$ $t_{av} = \frac{t_1 + t_2 + t_3}{3} =$ | $v_{av4} =$ |
| 100 cm | $h_5 =$ | $t_1 =$ $t_2 =$ $t_3 =$ $t_{av} = \frac{t_1 + t_2 + t_3}{3} =$ | $v_{av5} =$ |
| 120 cm | $h_6 =$ | $t_1 =$ $t_2 =$ $t_3 =$ $t_{av} = \frac{t_1 + t_2 + t_3}{3} =$ | $v_{av6} =$ |

Table 4.1: Repeated measurement at each position leading to v_{avg} needed for evaluating KE

| Position s (cm) | y_i vertical (mm) | $h_i = y_i - y_6$ (mm) | height h_i (m) | $PE = mgh$ (J) | v_{avg} (m/s) | $KE = \frac{mv^2}{2}$ (J) | $ME = PE + KE$ (J) |
|----------------------|------------------------|---------------------------|---------------------|-------------------|--------------------|------------------------------|-----------------------|
| 20 | | | | | | | |
| 40 | | | | | | | |
| 60 | | | | | | | |
| 80 | | | | | | | |
| 100 | | | | | | | |
| 120 | | | | | | | |

Table 4.2: Data leading to KE, PE and their sum

Part III

The World of Electricity

Chapter 5

Lab 5: Electricity

Name _____ Section _____ Date _____

5.1 Objectives

To study some consequences of three fundamental principles of electricity:

1. Like charges repel, and unlike charges attract.
2. Electrons, which are negatively charged, move freely in a conductor, but cannot move through an insulator. Positive charges cannot move in solids, even if they are conductors.
3. A *neutral* material has as much positive charge as negative charge.

You should be able to explain all the observations you make in this lab with these ideas.

5.2 Electrostatics

- Figure 5.1 shows the electroscope. The essential feature is that two gold-foil leaves hang from a metal rod connected to a conducting ball. To begin with, make sure the leaves hang freely.
- Rub a piece of acetate (the clear cylindrical plastic rod) with a dry and clean piece of paper and bring the acetate close to the silver conducting ball on the top of the electroscope. Then, touch the conducting ball on top of the electroscope with the acetate. Next, touch the ball with your hand. Next, repeat the process with the aluminum bar.
- Note: Sometimes the gold foil leaves will stick to themselves or to the glass plates on the front of the electroscope. This is undesirable, and you should try to avoid it when you do the experiments.




Figure 5.1: The electroscope

Question #:1 Describe carefully what you observe as you perform the above experiments.

Question #:2 The acetate is *positively charged* after being rubbed with the paper. Use this fact to explain your observations. Specifically, explain

1. how electrons move in the electroscope as you bring the acetate *close* (but not touching) to the electroscope. Make a sketch of the electroscope and acetate strip showing regions of positive and negative charge.
2. how electrons move when you touch the acetate to the conducting ball. Again, make a sketch of the electroscope and acetate strip showing how charge flows, and what the final charge distribution looks like after the acetate strip has been taken away.
3. why the gold-foil leaves move in each case.
4. what happens to the charge on the electroscope when you touch it with your hand.

Question #:3 The gold foil is connected to the ball by a metal rod, and this experiment shows that charge can be transferred through the rod to the gold foil. Why does the experiment not work when you touch the electroscope case with the acetate instead? (Try it).

- Ground the electroscope by touching the silver ball with your hand. The gold foil should return to its neutral “resting” position.
- Now, rub the vinyl strip (the grey rod) with the paper and touch the strip to the silver ball on the electroscope.
- Then, rub the acetate strip with the paper and bring it close to the ball on the electroscope.
- Finally, touch the acetate to the silver ball.

Question #:4 Describe carefully what you observe in the above steps.

Question #:5 What do your observations tell you about the charge on the vinyl strip? Give clear reasons. Make sketches showing (a) how charge moves when you bring the vinyl strip close to the electroscope and (b) how charge moves when you touch the electroscope with the vinyl strip.

- Ground the electroscope by touching the silver ball with your hand.
- Charge the electroscope with either the acetate or vinyl strip.
- Several pith-balls, which can be suspended by a nylon thread, are provided. Ground a pith ball by touching it with your hand. Then hold the pith-ball by the nylon thread, so it can swing freely and bring it *slowly* closer to the electroscope (without swinging it). Look carefully for the effects.

Question #:6 Describe carefully what you observe happening before, during, and after contact occurs between the pith-ball and the electroscope. Things happen quickly, and can be hard to see, so you may want to repeat this experiment several times. Also, note whether the gold-foil moves.

Question #:7 Explain your observations in terms of the motion of electrons. Note that the pith-ball is a conductor, and that your answer to question 3 is important here. Divide your explanation into three categories:

1. Before the pith-ball makes contact with the electroscope.
2. During contact.
3. After the pith-ball makes contact with the electroscope.

5.3 The Concept of “Ground”

The concept of “ground” is extremely important in electricity. It refers to the fact that, given the chance, excess electrical charge will almost always flow into the earth. The next few questions elaborate on this point.

Think of the earth as a giant conducting sphere (yes, dirt conducts electricity, though not particularly well). Use the principles outlined in the Objectives and think about your answer to the last prelab question to answer the following questions:

Question #:8 Why does touching the electroscope with your hand discharge it completely, but touching it with a pith-ball only discharge it a little bit?

Question #:9 Given the choice, why would excess charge on the electroscope prefer to flow into the earth, rather than into (for example) your body?

Question #:10 All modern electrical appliances are “grounded”. This means that the metal exterior of your toaster is connected to the earth by a wire. Why? (Hint: This is a safety feature. Think about what would happen if a wire was broken and touched the side of the toaster.)

5.4 Conductors and Insulators:

- Most materials fall into one of two categories: either they are electrical *conductors* or electrical *insulators*. In fact, almost nothing is a perfect conductor or insulator, and most materials fall in between.
- We can describe how good a conductor a material is using the concept of *electrical resistance*. If the resistance is low, the material is a good conductor. If the resistance is high, the material is a poor conductor. A perfect conductor has no resistance and a perfect insulator has infinite resistance.
- Fill out the table below using the instructions described afterwards:

| Material | Resistance |
|--------------|------------|
| aluminum rod | |
| plastic rod | |
| glass rod | |
| wooden block | |
| human hand | |
| copper wire | |
| air | |




Figure 5.2: A simple electrical circuit.

- Charge the electroscope with either the vinyl or acetate strip. Then discharge the electroscope by touching the conducting ball with the aluminum bar. Notice (roughly) how long it takes for the electroscope to discharge, and use this to determine the resistance of the bar.
- In the second column, indicate whether the electrical resistance is low, medium, or high.
- Repeat the process for each material provided, as well as any others you want to include.

Question #:11 When electricity was first introduced into homes, many people were nervous about the fact that electrical sockets didn't have plugs. They were worried that if nothing was plugged into the sockets, electricity would leak out of the holes into the room. Why doesn't electricity leak out of electrical sockets?

5.5 Simple Circuits:

So far, you have established that electrons can flow through conductors, and that they will do so (a) to move away from areas of excess negative charge (b) to move towards areas of excess positive charge. This flow is called an electrical current. Electrical currents are useful to us because we can make the electrons do *work* while they move.

You can think of a power supply as a device which builds up excess positive and negative charges on its two terminals. If some conducting path is available, electrons will flow between the terminals because of the forces between the charges. In the next experiments, the electrons will do work illuminating a lightbulb as they flow.

The important difference between a power supply (like a battery) and the electroscope is that the power supply replenishes the charge on its terminals whenever it is depleted.

- Construct the simple circuit shown in Fig. 5.2. The “-” sign on the power supply indicates that there is a surplus of negative charge at that terminal.




Figure 5.3: A more complicated electrical circuit.

- When the power supply is turned on, the lightbulb is illuminated, indicating that electrons are flowing through the circuit.

Question #:12 Draw a sketch like Fig. 5.2 and show the direction electrons flow through the wire. Why do they flow in this direction? (Hint: Think about the forces on the electrons.)

Question #:13 Describe what happens if you insert a piece of aluminum into the circuit? (See Fig. 5.3) What about a piece of wood? What about some of the other materials you looked at earlier?

5.6 Summary & Conclusions

Summarize briefly what this lab demonstrated. Think about the following questions: What makes electrical charges move? What factors affect how much charge moves? Why is the concept of resistance useful for describing this?

Question #:14 Different materials have different electrical properties. For example, glass is an insulator, and aluminum is a conductor. Is it (a) because they contain different numbers of electrons or (b) because the electrons move more freely in some materials than others? Is there any evidence from this lab you can use to support your answer?

Part IV

Within the Atom

Chapter 6

Lab 6: Atomic Spectra

Name _____ Section _____ Date _____

Caution! *The fluorescent gas tubes used in this lab are delicate, and become very hot during use. Be careful when you handle them. The fluorescent gas tubes in this lab may contain mercury. Mercury is poisonous. If you break a gas tube, notify your TA immediately.*

6.1 Objectives

1. To compare the spectra of a tungsten filament, a fluorescent light tube, and hydrogen tube with each other.
2. To study two different theories of light emission.

6.2 The spectroscope and the visible spectrum

- The eyepiece of the spectroscope contains a *diffraction grating* which breaks light up into its constituent colors. Use the spectroscope to look at an *incandescent* light bulb. The colors in the light emitted by the bulb will appear to be superimposed on a scale, as shown in Fig. 6.1(b). The scale tells you the wavelength of the light in hundreds of nanometers (nm). Recall that 10^9 nm is 1 m. Also recall that for multiplying scientific notation, the exponents belonging to the power of ten add, and for dividing the exponents subtract, so that $\frac{10^5}{10^{-3}} = 10^8$ since $5 - (-3) = 8$.

If you have a scientific calculator, you might have a button with an E on it, which denotes scientific notation. With this notation $4E5 = 4 \times 10^5$. Do not confuse this with the button which says e, which stands for a number approximately 2.72.




Figure 6.1: Spectroscope and spectroscopy scale (for the old spectroscopes no longer used in the lab).

Question #:1 Make another table like the one shown below. This one does not have enough room.

Table 1: Spectrum of the Incandescent Bulb

| Color | Approx. Range of Wavelengths (nm) | |
|-------|-----------------------------------|--|
| | | |

Use this table to record what you see when you look at the spectrum of the incandescent bulb. In the table, indicate the colors you see, (use 6-7 names for basic colors that you see), and the range of wavelengths you see them in.

Question #:2 What color has the longest wavelength? What color has the shortest wavelength? What colors are the brightest and dimmest? Wavelength is measured by the placement of the color on the scale, not the length the color seems to exist for.

Question #:3 Why do you think the spectrum cuts off beyond certain wavelengths? Does it mean that the lightbulb only produces the colors you see?

Question #:4 Light with wavelengths shorter than about 400 nm is called “ultraviolet”, and light with wavelengths longer than about 650 nm is called “infrared”. Why is this?

Question #:5 The relationship between wavelength and frequency is

$$f = \frac{c}{\lambda} \quad \text{or} \quad \lambda = \frac{c}{f}$$

where c is the speed of light, λ (the Greek letter “lambda”) is the wavelength *measured in metres* and f is the frequency in Hertz (Hz). The speed of light is $c = 3 \times 10^8$ m/s. What is a typical frequency of red light? What is a typical frequency of violet light? Show your calculation.

Question #:6 What aspects of spectrum of the light bulb does Hypothesis 1 in the pre-lab describe correctly? Are there any predictions it makes which are incorrect?

6.3 Spectra of Gases

- Use the spectroscope to study the spectrum of one of the overhead fluorescent lights.

- Now use the spectroscope to study the spectrum of hydrogen.

Question #:7 Describe the two spectra carefully. How do they compare with each other and with the spectrum of the incandescent light bulb?

Question #:8 Can your observations be explained in terms of random motion of electrons in the gas tubes? Why or why not? (Think carefully about this question when you write your conclusions at the end of the lab).

6.4 Measuring Spectra

6.4.1 Calibrating the Spectroscope

- To see how accurate the scale in the spectroscope is, it can be compared to a known source.
- Look at the hydrogen tube with the spectroscope and measure the wavelengths of the lines. Notice that you can adjust their width using the screw at the end of the spectroscope (don't force the screw). Wider lines are easier to see, but more exact measurements come from thinner ones.
- Complete the table below:

Calibration Data

| Measured λ (nm) | Actual λ (from chart) (nm) | Actual–Measured (correction) (nm) |
|----------------------------|--|---|
| | | |

The *correction* is the difference between the actual λ and the measured λ .

- Be sure to note whether the measurements are too low or too high.
- Be sure to note whether the correction is biggest at red wavelengths or at violet wavelengths.

6.4.2 Measuring Unknown Gases

- There are several unknown tubes provided.

Question #:9 Compare their spectra with the wall chart. What are the unknown gases? Match each one up to a substance. Distinguish the gasses either by numbering shown on the boxes, or by what color they seem to glow (without a spectroscope).

6.5 Bohr's atomic model and Plack's constant

Neils Bohr proposed a *quantum theory* of light emission from atoms. You will learn more about this theory in class. For now, all you need to know is:

Hypothesis 2 *According to Bohr's theory, the gas tube is produces light at only certain specific frequencies. The allowed frequencies are:*

$$f = Z \times (3.29 \times 10^{15}) \left[\frac{1}{n^2} - \frac{1}{m^2} \right]$$

where Z is the number of protons in the nucleus of the gas atom ($Z = 1$ for hydrogen, $Z = 2$ for helium, and so on). The numbers n and m can be $1, 2, 3 \dots$ and we only consider $m > n$.

Question #:10 Complete the table below using Bohr's formula for the predicted spectrum of hydrogen. Note that you are only calculating a small fraction of the possible frequencies.

| m | n | f (Hz) | λ (m) | λ (nm) |
|-----|-----|-------------|------------------|-------------------|
| 3 | 1 | | | |
| 3 | 2 | | | |
| 4 | 2 | | | |
| 5 | 2 | | | |

Question #:11 Which of the wavelengths you just calculated are visible?

Question #:12 How well do your calculated lines compare with the spectrum you measured.

6.6 Summary & Conclusions

Go back and look at the objectives for this lab. Write a brief summary of your findings, making sure you address the issues raised in the objectives. Remember: you studied two hypotheses for light emission. How well did they work? How are they different? What things do they get right? What things do they get wrong? Could either of the theories be used to explain both the tungsten filament *and* the hydrogen gas tube?

Question #:13 [Bonus Question] Now consider helium ($Z = 2$). Calculate the wavelength of light produced in the transition of an electron from $m = 4$ to $n = 3$. Show your calculations clearly. How does the calculated wavelength compare with the spectrum on the wall chart.

Chapter 7

Lab 7: Radioactivity

Name _____ Section _____ Date _____

Caution: *Liquid nitrogen is extremely cold and can cause severe damage if it makes prolonged contact with skin. Spills are especially dangerous if the nitrogen soaks into clothing.*

7.1 Objective

1. To study background radiation.
2. To determine the kinds of radiation emitted from a lead-210 sample.
3. To determine the levels of the various kinds of radiation emitted by the lead-210 sample.

7.2 Radiation

7.2.1 Kinds of radiation

- Alpha (α) particles: Very large nuclei may spontaneously emit an alpha particle. For example, Americium, which is near the bottom of the periodic table, decays this way. The α -particle is a helium (He) nucleus consisting of two protons and two neutrons. It is therefore *positively* charged. α -particles, which are relatively heavy, will produce straight dense trails in a cloud chamber. α -particles have a range of less than 8 cm in air, and can be stopped by a piece of paper.
- Beta (β) particles: If a nucleus has too many neutrons the most likely form of radioactive decay will be the emission of an electron from the nucleus. These high-energy electrons are known as beta particles. The β -particle is formed by the instant transformation of a neutron into a proton

plus an energetic electron which then escapes from the nucleus. β -particles are light and leave wispy, irregular trails. β -particles have a typical range of 10–10000 cm in air, and 0.01-10 cm in water, depending on how fast they are travelling. β -particles are *negatively* charged.

- Gamma (γ) rays: After each of the previous types of radioactive decay the new nucleus will have an excess of energy and this is usually released by the emission of one or more gamma-rays . γ -rays are electromagnetic radiation similar to radio waves, visible light, and X-rays, except that they have a much higher frequency (or shorter wavelength). γ -rays will penetrate to great depths in materials, and no amount of absorber will completely stop all of the gamma radiation . What is usually done in practice is to use sufficient thickness of an absorber to reduce the radiation level to an acceptable value. γ -rays have *no* charge, and are difficult to see in a cloud chamber.
- Cosmic rays: When there is no radiation source, cosmic rays may enter the chamber, producing thin misty trails. Cosmic rays are extremely energetic particles, primarily protons, which originate in the sun, other stars and some of the violent events which occur in space. The cosmic ray particles interact with the upper atmosphere of the earth and produce showers of lower energy particles. Many of these lower energy particles are absorbed by the earths' atmosphere as they travel down to the surface. At sea level the cosmic radiation is composed mainly of muons, with some gamma-rays, neutrons and electrons. As cosmic rays bounce off other particles in the cloud chamber, they produce “curly” or jagged tracks.

7.3 The Geiger Counter

The Geiger counter is a sensitive tool for measuring β - and γ -rays (α -rays cannot penetrate the glass case of the detection tube). The Geiger counters used in this lab emit an audible “click” when they detect a radioactive particle. These particular Geiger counters only operate when you push the button on their front.

Radiation is detected in a sensitive “detection tube” which is, in this case, the gold-coloured tube embedded in the side of the Geiger counter. Measure the dimensions of the detection tube and calculate the cross sectional area of the tube in cm^2 .

Measure the background radiation of the room. State how you measured the radiation level, and give the value. The simplest measurement is to count the number of clicks (or *counts*) over some time interval. Convert the amount of radiation you measure into the amount of radiation per cm^2 per second (ie. $\text{counts}/\text{cm}^2\text{s}$).

Estimate your cross-sectional surface area in cm^2 (describe how you make this estimate). From this, how much radiation is passing through your body per second?

Now take the Geiger counter outside and measure the background radiation. How many counts/ cm^2s do you measure outside? Is it different from inside? Why do you think this is?

Based on differences between radiation levels inside and outside, where do you think the background radiation is coming from?

Hold the radioactive source a few cm away from the detection tube (the gold-coloured tube) in the Geiger counter. How does the radiation level compare with the background radiation? How far do you have to move away from the source before the radiation is indistinguishable from the background?

Give two reasons why the radiation measured by the Geiger counter drops as you move the source away.

- Hold the radioactive source a few cm away from the detection tube (measure and the distance). Record the distance. Now measure and record the level of radiation from the source, using the same method as you used to measure the background radiation.

- Now hold the source *behind* the Geiger counter, so that the radiation must travel through the black case to reach the detection tube. Make sure the distance to the detection tube is the same as above. Now measure and record the radiation level. Don't forget to mention the time interval you measure over.
- Make a table like the one below

| Source position | # of counts | # of counts (background subtracted) | Source radiation level (counts/cm ² s) |
|-----------------|-------------|-------------------------------------|---|
| In Front | | | |
| Behind | | | |

- In the first column, record the total number of counts you detect. In the second column, subtract the number of counts which can be attributed to background radiation. In the final column, determine the radiation level measured by the detector.

When the detection tube is visible to the radioactive source, both β - and γ -rays are detected (along with the background), but when the source is behind the Geiger counter β -rays are screened by the black case and only γ -rays (and the background radiation) are detected. What are the levels of β - and γ -radiation you measured from the source?

Obviously, the number you get in the previous question is going to depend on how far the detector is from the source. You can estimate the *total* radiation of each kind (in units of counts/second) emitted by the source by assuming that the radiation is spread out evenly over the surface of a sphere. If the detector was d cm from the source in your experiment, then the total radiation passing through an imaginary spherical shell enclosing the source is

$$\text{total radiation} \left[\frac{\text{counts}}{\text{second}} \right] = \text{measured radiation} \left[\frac{\text{counts}}{\text{cm}^2 \text{ second}} \right] \times 4\pi d^2 [\text{cm}^2].$$

Estimate the total radiation levels of β - and γ -rays produced by your source.

By counting the number of “clicks”, you learn something about radiation levels in the room. This isn’t the whole story though. Counting “clicks” doesn’t say anything, for example, about whether the radiation is dangerous. What other information would be useful to know about the radiation? Hint: You know that radiation is just another name for “fast-moving particles”. If the particles were bigger (for example, baseball sized) what kinds of things would determine whether they were dangerous?

Write down the equation for the β -decay of ^{210}Pb (Lead-210). What is the isotope which is produced in this reaction?

The β -particle is a fast moving electron and has kinetic energy. The γ -ray carries radiant energy. Where does this energy come from?

7.4 The Cloud Chamber

7.4.1 Operating the cloud chamber

Making the cloud chamber work properly is not easy. You will need to spend time experimenting to establish the best operating conditions. If you are having trouble seeing anything, be patient.

- Place the green blotter paper and black cardboard disk into the cloud chamber. Make sure the hole in the side of the cloud chamber is not blocked by the blotter paper.
- Soak the green blotter paper and black cardboard disk with alcohol. Be generous with the alcohol, but do not add so much alcohol that a pool forms at the bottom. Put the lid on the cloud chamber.

- A radioactive source is provided, mounted on a cork. Insert the cork into the hole in the side of the chamber, so that the source is inside.
- Place the aluminum stand in the center of the styrofoam dish. Fill the dish about half-way with liquid nitrogen. Liquid nitrogen will be given to you by the TA.
- Place the cloud chamber on the aluminum stand. Illuminate the interior of the cloud chamber through the side window with the light source.
- Now watch the cloud chamber carefully. Nothing will happen for several minutes. As the chamber cools, you will start to see tracks left by passing radiation. The number of tracks you see will depend on how well you have optimized operating conditions.

7.4.2 Optimizing the operating conditions

To optimize the chamber properly, you should know a little bit about how it works. When a subatomic particle (radiation) travels through the chamber, it collides with air molecules, producing free ions. Alcohol vapor in the chamber condenses around these free ions, forming droplets. The droplets are what form the trail. Ideally,

1. the alcohol-soaked blotting paper should be warm to produce lots of vapor,
2. one other surface (in this case the base) should be cold.

The cold surface produces a layer of cold air. Alcohol vapor in the cold air layer wants to condense to form clouds, but needs something—in this case, the free ions—to “seed” or “nucleate” the condensation.

What can go wrong?

- The chamber is too warm. In this case, the alcohol vapor doesn’t want to condense. No tracks are seen.
- The chamber is too cold. In this case, the alcohol vapor condenses too easily, and forms fog. If this happens, take the cloud chamber off the aluminum block for awhile.
- The blotting paper is too cold. In this case, no alcohol vapor is produced. Again, take the cloud chamber off the aluminum block.
- *The most common problem is that the cloud chamber becomes too cold. Remember that the aluminum block takes a long time to cool down, but will also keep the chamber cold for a long time. It is not necessary to add more liquid nitrogen immediately after you run out!*
- Once you have learned how to operate the cloud chamber, answer the following questions:

Question #:1 Describe the different kinds of radiation tracks you see. Where do the tracks come from? Do they all come from the same place? Which kinds do you see most often?

Question #:2 Read the descriptions of the different kinds of radiation at the beginning of this lab. Based on your observations, what kinds of radiation are you observing in the cloud chamber? Is this consistent with your Geiger counter measurements?

Question #:3 Compare the radiation levels you see in your cloud chamber with your earlier estimate of the radiation produced by the source. Do you expect these numbers to be the same? Why?

Question #:4 Take the source out of the cloud chamber. Describe the radiation tracks you observe. What kind of radiation are you seeing?

7.5 Summary & Conclusions

Summarize your findings, remembering to address issues raised in the objectives.

